



ambetter.[®]

Marketplace 2017 – P4P Program

HEDIS Quick Reference Guide



Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis

The percentage of adults 18–64 years of age with a diagnosis of acute bronchitis who were not dispensed an antibiotic prescription.

The measure is reported as an inverted rate $[1 - (\text{numerator}/\text{eligible population})]$. A higher rate indicates appropriate treatment of adults with acute bronchitis (i.e., the proportion for whom antibiotics were not prescribed).

Proportion of Days Covered

The percentage of patients 18 years and older who met the proportion of days covered (PDC) threshold of 80% during the measurement period. Rates are reported for the: Diabetes All Class, Renin Angiotensin System (RAS) Antagonists, and Statins.

Use of Imaging Studies for Low Back Pain

The percentage of members with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.

The measure is reported as an inverted rate $[1 - (\text{numerator}/\text{eligible population})]$. A higher score indicates appropriate treatment of low back pain (i.e., the proportion for whom imaging studies did not occur).

Annual Monitoring of People with Persistent Medications

The percentage of members 18 years of age and older who received at least 180 treatment days of ambulatory medication therapy for a select therapeutic agent during the measurement year and at least one therapeutic monitoring event for the therapeutic agent in the measurement year.

- **Angiotensin** converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARB)
 - At least one serum potassium and a serum creatinine therapeutic monitoring test in the measurement year
- **Digoxin**
 - At least one serum potassium, at least one serum creatinine *and* at least one serum digoxin therapeutic monitoring test in the measurement year
- **Diuretics**
 - At least one serum potassium *and* a serum creatinine therapeutic monitoring test in the measurement year

Medical Management of People with Asthma

The percentage of members 5–64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period.

- The percentage of members who remained on an asthma controller medication for at least 75% of their treatment period.



Cervical Cancer Screening

Measure evaluates the percentage of women ages 21–64 who were screened for cervical cancer using either of the following criteria:

- 1) Cervical cytology performed every 3 years for women ages 21–64
- 2) Cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years (must occur within 4 days of each other) for women ages 30–64.

Women who have had a hysterectomy without a residual cervix are exempt from this measure.

Cervical Cytology plus one HPV code Ages 30-64		Cervical Cytology Codes Ages 21-64	
<u>CPT</u> 87620, 87621, 87622, 87624, 87625	<u>HCPCS</u> G0476	<u>CPT</u> 88141-88143, 88147, 88148, 88150, 88152- 88154, 88164-88167, 88174, 88175	<u>HCPCS</u> G0123, G0124, G0141, G0143- G0145, G0147, G0148, P3000, P3001, Q0091

Breast Cancer Screening

Measure evaluates the percentage of women ages 50–74 who had a mammogram at least once in the past two years. Women who have had a bilateral mastectomy are exempt from this measure.

<u>CPT</u>	<u>HCPCS</u>
77055-77057	G0202, G0204, G0206

Colorectal Cancer Screening

Measure evaluates the percentage of members ages 50-75 who had at least one appropriate screening for colorectal cancer in the past year. Appropriate screening is FOBT in 2017, sigmoidoscopy in the last 5 years or colonoscopy in last 10 years. Patients who have a history of colon cancer or who have had a total colectomy are exempt from this measure.

FOBT		Flexible Sigmoidoscopy		Colonoscopy	
<u>CPT</u> 81528, 82270, 82274	<u>HCPCS</u> G0328, G0464	<u>CPT</u> 45330-45335, 45337-45342, 45346-45347, 45349-45350	<u>HCPCS</u> G0104	<u>CPT</u> 44388-44394, 44397, 45355, 45378-45387, 45391, 45392, 74263	<u>HCPCS</u> G0105, G0121



Diabetes Care (Comprehensive)

Measure demonstrates the percentage of members ages 18-75 with diabetes (types 1 & 2) who were compliant in the following sub measures:

HbA1c Test: is completed at least once per year (includes rapid A1c).

CPT

83036, 83037

CPT II

3044F, 3045F, 3046F

HCPCS

Nephropathy Screening Test: is performed at least once per year. A member who is on ACE/ARBs or has nephropathy is compliant for this sub measure.

CPT

82042, 82043, 82044, 81000,
81001, 81002, 81003, 81005,
84156

CPT II

3060F, 3061F, 3062F

HCPCS

Eye Exam: a retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) is complete every year OR a negative retinal exam (no evidence of retinopathy) by an eye care professional in the year prior. CPT II code 3072F reflects a dilated retinal exam negative for retinopathy.

CPT

67028, 67030, 67031, 67036,
67039-67043, 67101, 67105,
67107, 67108, 67110, 67112,
67113, 67121, 67141, 67145,
67208, 67210, 67218, 67220,
67221, 67227, 67228, 92002,
92004, 92012, 92014, 92018,
92019, 92134, 92225-92228,
92230, 92235, 92240, 92250,
92260, 99203-92205, 99213-99215,
99242-99245, 92134

CPT II

2022F, 2024F, 2026F, 3072F

HCPCS

S0620, S0621, S3000