

Clinical Policy: Inhaled Nitric Oxide

Reference Number: CP.MP.87

Effective Date: 08/13

Last Review Date: 09/17

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Inhaled Nitric Oxide (iNO) is a selective pulmonary vasodilator in which its mechanism of action results in smooth muscle relaxation. Several studies have suggested that iNO improves oxygenation, particularly in trials of term and near-term neonates with hypoxic respiratory failure. iNO has been shown to reduce the need for ECMO (extracorporeal membrane oxygenation) without increasing neurodevelopmental, behavioral, or medical abnormalities at 2 years of age.

Policy/Criteria

- I. It is the policy of health plans affiliated with Centene Corporation[®] that iNO therapy is **medically necessary** for the following indications:
 - A. *Initiation* of therapy
 1. Hypoxic respiratory failure in newborns ≥ 34 weeks gestational age at birth with all:
 - a. Pulmonary artery hypertension (PAH) diagnosed by echocardiogram that excluded congenital heart disease (CHD), *and*
 - b. Conventional therapies such as mechanical ventilation, administration of high concentrations of oxygen (80-100%), high-frequency ventilation, induction of alkalosis, neuromuscular blockade, and sedation have failed or are expected to fail; *and*
 - c. Oxygen index (OI) ≥ 25 . The OI is calculated as the mean airway pressure divided by the partial pressure of arterial oxygen times 100; *and*
 - d. Response seen with administration of up to 40 ppm trial of iNO (defined as a PaO₂ increase ≥ 20 mm Hg or a 20% decrease in OI.
 2. Perioperative management of pulmonary hypertension in infants and children with CHD, must meet all:
 - a. iNO therapy for vasodilation is used in response to cardiac bypass surgery to repair a congenital heart defect that is causing PAH, *and*
 - b. iNO is delivered directly to the lungs via endotracheal tube.
 - B. Continuation of iNO therapy in newborns
 1. Member continues to require iNO as evidenced by a continued O₂ requirement of 80-100%, or
 2. A weaning protocol is being initiated after a 4-6 hour period of stability indicated by O₂ requirement decreased to 60-80% or OI ≤ 10 ; *and*
 - a. The iNO has been used for 1 week or less
- II. It is the policy of health plans affiliated with Centene Corporate that iNO is **not medically necessary** for any other indications such as preterm infants < 34 weeks gestation, acute bronchiolitis, bronchopulmonary dysplasia (BPD), congenital diaphragmatic hernia (CDH),

adult respiratory distress syndrome or acute lung injury, treatment in adults with positive vaso-reactivity testing, post-op cardiac surgery in adults, and vaso-occlusive crises in members with sickle cell disease because safety and effectiveness have not been established.

Treatment Regimen

The American Academy of Pediatrics (AAP) recommends that iNO should only be administered according to a formal protocol that has been approved by the Food and Drug Administration (FDA) and the institutional review board and with informed consent. Use should be limited to sites with multisystem support, including on-site ECMO capability. If ECMO is not available on-site, a timely transfer needs to be arranged to a collaborating ECMO center without interruption of iNO therapy.

Since no one standard protocol has been issued for iNO treatment, the following is one guideline to assist in determining appropriate initiation and continuation of treatment. The recommended starting dose of iNO for term infants is 20ppm. A positive response generally occurs in less than 30 minutes with a PaO₂ increase ≥ 20 mmHg (or 20% decrease in OI). If there is no response, the dose may be increased up to 40 ppm. In premature infants, the initial dose used in studies was 10 ppm with an increase up to 20 ppm in non-responders. Doses of up to 80 ppm have been used, but the potential for increasing toxicity without additional benefits occurs at doses greater than 40 ppm.

Per Peliowski, weaning can occur following improvement in oxygenation and after a 4 to 6 hour period of stability, during which the inspired oxygen concentration is decreased to 60% to 80%, or the OI falls to ≤ 10 . At 4-6 hour intervals, the dose can be decreased by 50%, as long as the OI remains ≤ 10 . When stability is maintained at iNO dose of 5 ppm, weaning should occur by 1 ppm every 4 hours and discontinued at 1 ppm if oxygenation status remains with $< 60\%$ oxygen with PaO₂ consistently > 50 mmHg. If deterioration occurs during or after weaning occurs, the dose should be increased to the previous level or iNO restarted. Once the infant stabilizes again, weaning should occur more slowly, taking place over a 24 to 48 hour period.

In general, patients who responded to iNO therapy typically require treatment for 3-4 days, with randomized trials demonstrating 90% of treated infants were off iNO therapy within one week of initiation. Patients should be monitored for potential toxic effects by measuring the serum methemoglobin concentration, levels of nitrogen dioxide at the airway opening, and ambient air contamination. Decreased platelet aggregation, increased risk of bleeding and surfactant dysfunction can also occur from iNO toxicity.

Background

A large and well-designed multicenter trial was conducted by the Neonatal Research Network in 235 infants with gestational age ≥ 34 weeks who had severe hypoxic respiratory failure (OI ≥ 25) and did not have CDH. Infants were randomly assigned to iNO or control (100% oxygen). Fewer infants in the treatment group died within 120 days or received ECMO therapy, (46% versus 64%; relative risk 0.72, 95% CI 0.57-0.91) compared to control. This difference was entirely due to decreased requirement for ECMO (39% versus 54%); there was no difference between groups in mortality.

In a systemic review by the Cochrane database, similar findings of fewer requirements for ECMO and no difference in mortality were noted. Fourteen randomized trials were found in term or near term infants with hypoxia. iNO improved oxygenation in approximately 50% of the treated infants. Within 30 to 60 minutes of beginning therapy, PaO₂ increased by a mean of 53 mmHg and OI decreased by a mean of 15.1. Outcome did not appear to be affected by whether infants had echocardiographic evidence of persistent pulmonary hypertension. No benefit was noted in those with CDH, indeed there is a suggestion that outcome was slightly worsened.

In preterm infants <35 weeks gestation, a systematic review by the Cochrane database found 14 randomized controlled trials of iNO. The authors concluded that iNO as a rescue therapy for the very ill ventilated preterm infant does not appear to be effective and may increase the risk of severe intraventricular hemorrhage. Later use to prevent BPD does not appear to be effective. Early routine use of iNO in mildly sick preterm infants may improve survival without BPD and decrease serious brain injury; further studies are needed to confirm these findings.

iNO has been well-studied in patients with acute lung injury and acute respiratory distress syndrome (ALI/ARDS). While iNO may improve oxygenation temporarily, it has not been shown to improve clinically important outcomes such as duration of mechanical ventilation, 28-day mortality or one-year survival. Furthermore, iNO does not improve oxygenation in all patients and the factors that may predict a good response are still uncertain.

In a systemic review by the Cochrane database, results from 5 randomized controlled trials found that iNO did not demonstrate any statistically significant effect on mortality and transiently improved oxygenation (up to 72 hours) in patients with hypoxemic respiratory failure. Further trials are needed to assess other clinically relevant end points and provide further stratification for treatment. In an updated Cochrane database review, the evidence was insufficient to support iNO in any category of critically ill adults and children with acute respiratory distress syndrome. Although, iNO results in a transient improvement in oxygenation, it does not reduce mortality and may be harmful, as it seems to increase renal impairment.²³

A Cochrane Summary for the use of iNO for pulmonary hypertension (PH) following surgery in infants and children with congenital heart disease found no benefit of it to assist in recovery. In the four randomized trials reviewed, there was no difference found in mortality or other outcomes reviewed. Due to the minimal data that was available, the authors found it difficult to draw valid conclusions regarding effectiveness and safety of this treatment in the select population. In a later study, iNO was effective in reducing the risk of development of PH crisis in PAH-CHD patients after cardiac repair in a placebo-controlled study¹⁶. Infants with PAH-CHD receiving iNO had fewer PH crises and shorter postoperative courses without concomitant side effects related to the medication.

Research on iNO use in adults with PH is limited to case reports and small case series, which leaves the impact of iNO on survival uncertain. It has been found to successfully stabilize a variety of acutely ill and hemodynamically compromised patients with severe PH, but the outcomes data are limited and thus cannot be considered standard of care. Acute vasodilator testing is the only well established and widely accepted use of iNO in patients with PAH.

CLINICAL POLICY
Inhaled Nitric Oxide

Patients with a positive vasoreactivity test are candidates for a trial of calcium channel blocker therapy

INO has numerous potential harms that must be considered when determining the risks and benefits of treatment. These potential harms include renal dysfunction, DNA strand breakage and base alterations which are potentially mutagenic, immunosuppression that could increase the risk of nosocomial infection, and a possible increase in methemoglobin and NO2 concentrations, which must be monitored frequently. Also, iNO may produce toxic free radicals; however, it is unknown if these are more harmful than ongoing exposure to high fractions of inspired oxygen.

Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2017, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
94799	Unlisted pulmonary service or procedure

HCPCS Codes	Description
N/A	

ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
I27.0	Primary pulmonary hypertension
I27.2	Other secondary pulmonary hypertension
P07.37	Preterm newborn, gestational age 34 completed weeks
P07.38	Preterm newborn, gestational age 35 completed weeks
P07.39	Preterm newborn, gestational age 36 completed weeks
P22.0	Respiratory distress syndrome of newborn
P28.5	Respiratory failure of newborn
P29.3	Persistent fetal circulation
Q21.0	Ventricular septal defect
Q21.2	Atrial septal defect

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed, Neonatologist reviewed	08/13	08/13
Split criteria into initiation and continuation of therapy	08/14	09/14

Reviews, Revisions, and Approvals	Date	Approval Date
Added Treatment Regimen section Neonatologist reviewed		
Literature researched and bibliography updated Converted to new template Neonatologist reviewed	09/15	09/15
Updated template and disclaimer language	03/16	
Added criteria for iNO therapy in perioperative management for PAH in pediatrics. Reviewed by Pediatric Cardiology, Adult Cardiology, and Neonatologist.	08/16	09/16
References reviewed and updated. ICD-10 and CPT codes added.	09/17	09/17

References

1. Adams JM, Stark AR. Persistent pulmonary hypertension of the newborn. In: UpToDate: Garcia-Prats JA (Ed), UpToDate, Waltham, MA. Update Feb 8, 2017 Accessed 09/11/17.
2. Afshari A, Brok J, Møller AM, Wetterslev J. Inhaled nitric oxide for acute respiratory distress syndrome (ARDS) and acute lung injury in children and adults. Cochrane Database of Systematic Reviews 2010, Issue 7. Art. No.: CD002787. DOI: 10.1002/14651858.CD002787.pub2
3. Allen MC, et al. Inhaled Nitric Oxide in Preterm Infants. Evidence Report/Technology Assessment No. 195. (Prepared by Johns Hopkins University Evidence-based Practice Center under Contract No. 290-2007-10061-1). AHRQ Publication No. 11-E001. Rockville, MD: Agency for Healthcare Research and Quality. October 2010.
4. American Academy of Pediatrics. Use of inhaled nitric oxide. Policy Statement. Pediatrics, August 2000, 106:2:344-345 (Reaffirmed December 2009)
5. Barrington KJ, Finer N. Review Inhaled nitric oxide for respiratory failure in preterm infants. Cochrane Database Syst Rev. 2010 Dec 8; (12):CD000509.
6. Bizzarro M, Gross I. Inhaled nitric oxide for the postoperative management of pulmonary hypertension in infants and children with congenital heart disease. Cochrane Database of Systematic Reviews 2005, Issue 4. Art. No.: CD005055. DOI: 10.1002/14651858.CD005055.pub2
7. Breuer J, Perin W, Gebhardt S, et al. . Inhaled nitric oxide treatment of children with pulmonary hypertension after cardiac surgery. Progress in Pediatric Cardiology, 1998, 2(9): 73-83.
8. Finer N, Barrington KJ. Nitric oxide for respiratory failure in infants born at or near term. Cochrane review abstract and plain language summary. The Cochrane Database of Systematic Reviews 2006 Oct 18;(4):CD000399.
9. Guthrie SO, et al. Initial dosing of inhaled nitric oxide in infants with hypoxic respiratory failure. J Perinatal. 2004 May;24(5): 290-4.
10. Granton J, Langleben D, Kutryk MJ, et al. Endothelial NO-Synthase Gene-Enhanced Progenitor Cell Therapy for Pulmonary Arterial Hypertension: the PHACeT Trial. Circ Res July 2015. 117:645-654 Available at: <http://circres.ahajournals.org/content/117/7/645.long>
11. Hopkins W, Rubin LJ. Treatment of pulmonary hypertension in adults. In: UpToDate, Mandel J (Ed), UpToDate, Waltham, MA. Update Apr 21 2017 Accessed 09/11/17.

CLINICAL POLICY
Inhaled Nitric Oxide

12. Ichinose F, Roberts JD Jr, Zapol WM. A selective pulmonary vasodilator: Current uses and therapeutic potential. *Circulation* 2004; 109: 3106-3111.
13. Inhaled nitric oxide for acute respiratory distress syndrome (ARDA) in adults. Hayes Health Technology Brief. March 4, 2015. Archived May 2017
14. Kumar P, Committee on Fetus and Newborn. Use of inhaled nitric oxide in preterm infants. *Pediatrics* 2014;133;164-170. Accessed at:
<http://pediatrics.aappublications.org/content/pediatrics/133/1/164.full.pdf>
15. Martin, R. Prevention and treatment of respiratory distress syndrome in preterm infants. In: UpToDate, Garcia-Prats JA (Ed), UpToDate, Waltham, MA. Update Aug 2, 2017. Accessed 09/11/17.
16. Miller OI, Tang SF, Keech A, et al. Inhaled nitric oxide and prevention of pulmonary hypertension after congenital heart surgery: a randomized double-blind study. *Lancet* 2000;356(9240):1464–1469.
17. Morales-Blanhir J, Sanos S, de Jover L, et al. , Clinical value of vasodilator test with inhaled nitric oxide for predicting long-term response to oral vasodilators in pulmonary hypertension, *Respir Med*, 2004: 98(3): 225.
<http://www.sciencedirect.com/science/article/pii/S0954611103003585>
18. Peliowski A, Canadian Paediatric Society, Fetus and Newborn Committee. Inhaled nitric oxide use in newborns. *Paediatr Child Health*. 2012 February; 17(2): 95-97.
<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3299357/>
19. Sokol J, Jacobs SE, Bohn D. Inhaled nitric oxide for acute hypoxemic respiratory failure in children and adults. Cochrane review abstract and plain language summary. *The Cochrane Database of Systematic Reviews* 2009.
20. Soll RF. Inhaled nitric oxide in the neonate. *Journal of Perinatology* (2009) 29, S63-S67.
21. Gebistorf F, Karam O, Wetterslev J, Afshari A. Inhaled nitric oxide for acute respiratory distress syndrome (ARDS) in children and adults. *Cochrane Database Syst Rev*. 2016 Jun 27;(6):CD002787
22. Siegel MD. Acute respiratory distress syndrome: Supportive care and oxygenation in adults. UpToDate, Waltham, MA. Update Aug 04, 2017. Accessed 09/11/17
23. Klingler JR, Inhaled nitric oxide in adults: Biology and indications for use. UpToDate, Waltham, MA. Update Aug 8, 2017. Accessed Sept 12, 17
24.

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

CLINICAL POLICY

Inhaled Nitric Oxide

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

Note: For Medicare members, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international

CLINICAL POLICY
Inhaled Nitric Oxide

copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.