

Clinical Policy: Glecaprevir/Pibrentasvir (Mavyret)

Reference Number: CP.PHAR.348

Effective Date: 09/17

Last Review Date: 09/17

Line of Business: Medicaid

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Glecaprevir and pibrentasvir (Mavyret™) are a fixed-dose combination of glecaprevir, a hepatitis C virus (HCV) NS3/4A protease inhibitor, and pibrentasvir, an HCV NS5A inhibitor.

FDA approved indication

Mavyret is indicated for the treatment of:

- Patients with chronic HCV genotype 1, 2, 3, 4, 5, or 6 infection without cirrhosis and with compensated cirrhosis (Child-Pugh A).
- Adult patients with genotype 1 infection, who previously have been treated with a regimen containing an HCV NS5A inhibitor or an NS3/4A protease inhibitor, but not both.

Policy/Criteria

Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with Centene Corporation® that Mavyret is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Chronic Hepatitis C Infection** (must meet all):

1. Diagnosis of chronic HCV infection as evidenced by detectable HCV RNA (ribonucleic acid) levels in the last 6 months;
2. Confirmed HCV genotype is one of the following (a, b, or c):
 - a. For treatment-naïve patients: genotypes 1, 2, 3, 4, 5, or 6;
 - b. For patients treatment-experienced with interferon (IFN)/pegylated-interferon (pegIFN), ribavirin (RBV), and/or sofosbuvir only: genotypes 1, 2, 3, 4, 5, or 6;
 - c. For patients treatment-experienced with either an NS5A inhibitor or an NS3/4A protease inhibitor: genotype 1 (*see Appendix D*);
3. Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist;
4. Age \geq 18 years;
5. Life expectancy \geq 12 months with HCV treatment;
6. Documented sobriety from alcohol and illicit IV drugs for \geq 6 months prior to starting therapy, if applicable;
7. Advanced liver disease defined as a or b:
 - a. Advanced fibrosis indicated by i or ii:
 - i. Liver biopsy showing a METAVIR score of F3 or equivalent (Knodell, Scheuer, Batts-Ludwig – F3; Ishak – F4/5);

CLINICAL POLICY**Glecaprevir/Pibrentasvir**

- ii. One serologic test and one radiologic test showing an equivalent score to METAVIR F3 per Appendix C;
 - b. Cirrhosis indicated by i, ii or iii:
 - i. Hepatocellular carcinoma (HCC) - and the HCC is amenable to resection, ablation or transplant;
 - ii. Liver biopsy showing a METAVIR score of F4 or equivalent (Knodell, Scheuer, Batts-Ludwig – F4; Ishak - F5/6);
 - iii. Both of the following:
 - a) One serologic test showing an equivalent score to METAVIR F4 per Appendix C;
 - b) One radiologic test showing an equivalent score to METAVIR F4 per Appendix C or other radiologic test showing evidence of cirrhosis (e.g., portal hypertension);
8. Prescribed regimen is consistent with an FDA or AASLD-IDSAs recommended regimen (*see Section V Dosage and Administration for reference*);
9. If cirrhosis is present, confirmation of Child-Pugh A status;
10. Member is hepatitis B virus (HBV) negative, or if positive, documentation that concurrent HBV infection is being treated (e.g., tenofovir alafenamide, adefovir, entecavir), unless contraindicated or clinically significant adverse effects are experienced (*see Appendix B*);
11. Member agrees to participate in a medication adherence program meeting both of the following components:
 - a. Medication adherence monitored by pharmacy claims data or member report;
 - b. Member's risk for non-adherence identified by adherence program or member/prescribing physician follow-up at least every 4 weeks;
12. Dose does not exceed glecaprevir 300 mg and pibrentasvir 120 mg (3 tablets) per day.

Approval duration: up to a total of 16 weeks*

(*Approved duration should be consistent with a regimen in Section V Dosage and Administration)

B. Other diagnoses/indications

1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

II. Continued Therapy**A. Chronic Hepatitis C Infection (must meet all):**

1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Mavyret for treatment of chronic HCV infection and has received this medication for at least 30 days;
2. Member is responding positively to therapy (e.g., decreased HCV RNA level, no unacceptable toxicity);
3. Dose does not exceed glecaprevir 300 mg and pibrentasvir 120 mg (3 tablets) per day.

Approval duration: up to a total of 16 weeks*

(*Approved duration should be consistent with a regimen in Section V Dosage and Administration)

CLINICAL POLICY

Glecaprevir/Pibrentasvir

B. Other diagnoses/indications

1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

III. Diagnoses/Indications for which coverage is NOT authorized:

- A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PHAR.57 or evidence of coverage documents;
- B.** Treatment-experienced patients with both NS3/4A protease inhibitor AND NS5A inhibitor, such as combination therapies including: Technivie, Viekira, and Zepatier.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

AASLD: American Association for the Study of Liver Diseases	IDSA: Infectious Diseases Society of America
DNA: deoxyribonucleic acid	IFN: interferon
HBeAg: hepatitis B virus envelope antigen	NS3/4A, NS5A/B: nonstructural protein
HBV: hepatitis B virus	pegIFN: pegylated interferon
HCC: hepatocellular carcinoma	PO: by mouth
HCV: hepatitis C virus	QD: once per day
FDA: Food and Drug Administration	RBV: ribavirin
FIB-4: Fibrosis-4 index	RNA: ribonucleic acid

Appendix B: General Information

- Hepatitis B Reactivation is a Black Box Warning for all direct-acting antiviral drugs for the treatment of HCV. The provider must provide either:
 - Documentation of absence of concurrent HBV infection as evidenced by laboratory values showing absence of hepatitis B virus envelope antigen (HBeAg) and HBV DNA (deoxyribonucleic acid);
 - Documentation that HBV co-infected patient may not be candidates for therapy as evidenced by one of the following:
 - Absence of HBeAg, HBV DNA less than 2,000 international units/mL, and alanine aminotransferase (ALT) level within 1 to 2 times the upper limit of normal;
 - HBeAg-positive and HBV DNA greater than 1,000,000 international units/mL and ALT level within 1 to 2 times the upper limit of normal;
 - Documentation that concurrent HBV infection is being treated (e.g., tenofovir alafenamide, adefovir, entecavir), unless contraindicated or clinically significant adverse effects are experienced.
- Due to higher rates of virologic failure and treatment-emergent drug resistance, the data do not support labeling for treatment of HCV genotype 1 infected patients who are both NS3/4A PI and NS5A inhibitor-experienced.

Appendix C: Approximate Scoring Equivalencies using METAVIR F3/F4 as Reference

Fibrosis/ Cirrhosis	Serologic Tests*				Radiologic Tests†		Liver Biopsy‡	
	Fibro Test	FIBRO Spect II	APRI	FI B-4	FibroScan (kPa)	MRE (kPa)	METAVIR	Ishak
Advanced fibrosis	≥0.59	≥42	>1.5	>3 .25	≥9.5	≥4.11	F3	F4-5
Cirrhosis	≥0.75	≥42	>1.5	>3 .25	≥12.0	≥4.71	F4	F5-6

*Serologic tests:

FibroTest (available through Quest as FibroTest or LabCorp as FibroSure)

FIBROSpect II (available through Prometheus Laboratory)

APRI (AST to platelet ratio index)

FIB-4 (Fibrosis-4 index: includes age, AST level, platelet count)

†Radiologic tests:

FibroScan (transient elastography)

MRE (magnetic resonance elastography)

‡Liver biopsy (histologic scoring systems):

METAVIR F3/F4 is equivalent to Knodell, Scheuer, and Batts-Ludwig F3/F4 and Ishak F4-5/F5-6

METAVIR fibrosis stages: F0 = no fibrosis; F1 = portal fibrosis without septa; F2 = few septa; F3 = numerous septa without cirrhosis; F4 = cirrhosis

Appendix D: Direct-Acting Antivirals for Treatment of HCV Infection

Brand Name	Drug Class				
	NS5A Inhibitor	Nucleotide Analog NS5B Polymerase Inhibitor	Non-Nucleoside NS5B Polymerase Inhibitor	NS3/4A Protease Inhibitor (PI)	CYP3A Inhibitor
Daklinza	Daclatasvir				
Epclusa*	Velpatasvir	Sofosbuvir			
Harvoni*	Ledipasvir	Sofosbuvir			
Olysio				Simeprevir	
Sovaldi		Sofosbuvir			
Technivie*	Ombitasvir			Paritaprevir	Ritonavir
Viekira XR/Pak*	Ombitasvir		Dasabuvir	Paritaprevir	Ritonavir
Zepatier*	Elbasvir			Grazoprevir	

*Combination drugs

V. Dosage and Administration

CLINICAL POLICY
Glecaprevir/Pibrentasvir

Indication	Dosing Regimen	Maximum Dose	Reference
Treatment-naïve Chronic hepatitis C (CHC) infection: Genotypes 1, 2, 3, 4, 5, or 6	Without cirrhosis: 3 tablets by mouth (PO) daily (QD) for 8 weeks With compensated cirrhosis: 3 tablets PO QD for 12 weeks	Glecaprevir 300 mg/pibrentasvir 120 mg (3 tablets) per day	FDA-approved labeling
Treatment-experienced with IFN/pegIFN + RBV +/- sofosbuvir CHC infection: Genotypes 1, 2, 4, 5, or 6	Without cirrhosis: 3 tablets PO QD for 8 weeks With compensated cirrhosis: 3 tablets PO QD for 12 weeks	Glecaprevir 300 mg/pibrentasvir 120 mg (3 tablets) per day	FDA-approved labeling
Treatment-experienced with IFN/pegIFN + RBV +/- sofosbuvir CHC infection: Genotype 3	Without cirrhosis or with compensated cirrhosis: 3 tablets PO QD for 16 weeks	Glecaprevir 300 mg/pibrentasvir 120 mg (3 tablets) per day	FDA-approved labeling
Treatment-experienced with NS5A inhibitor without prior NS3/4A protease inhibitor CHC infection: Genotype 1	Without cirrhosis or with compensated cirrhosis: 3 tablets PO QD for 16 weeks	Glecaprevir 300 mg/pibrentasvir 120 mg (3 tablets) per day	FDA-approved labeling
Treatment-experienced with NS3/4A protease inhibitor without prior NS5A inhibitor CHC infection: Genotype 1	Without cirrhosis or with compensated cirrhosis: 3 tablets PO QD for 12 weeks	Glecaprevir 300 mg/pibrentasvir 120 mg (3 tablets) per day	FDA-approved labeling

VI. Product Availability

Tablets: glecaprevir 100 mg and pibrentasvir 40 mg

VII. References

1. Mavyret Prescribing Information. North Chicago, IL: AbbVie Inc.; August 2017. Available at: www.mavyret.com. Accessed August 7, 2017.

Reviews, Revisions, and Approvals	Date	Approval Date
Policy created.	08/17	09/17

Important Reminder

CLINICAL POLICY

Glecaprevir/Pibrentasvir

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

CLINICAL POLICY
Glecaprevir/Pibrentasvir

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.