

## **Clinical Policy: Collagenase Clostridium Histolyticum (Xiaflex)**

Reference Number: CP.PHAR.82

Effective Date: 10.01.11

Last Review Date: 11.17

Line of Business: Medicaid

[Coding Implications](#)  
[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Collagenase clostridium histolyticum (Xiaflex<sup>®</sup>) is a combination of bacterial collagenases.

### **FDA Approved Indication(s)**

Xiaflex is indicated for the treatment of:

- Adult patients with Dupuytren's contracture with a palpable cord
- Adult men with Peyronie's disease with a palpable plaque and curvature deformity of at least 30 degrees at the start of therapy

### **Policy/Criteria**

Provider must submit documentation (which may include office chart notes and lab results) supporting that member has met all approval criteria

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Xiaflex is **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Dupuytren's Contracture** (must meet all):

1. Diagnosis of Dupuytren's contracture with a palpable cord and both (a and b):
  - a. Positive "table top test" (inability to simultaneously place affected finger and palm flat against a table top);
  - b. Flexion contracture  $\geq 20$  degrees in metacarpophalangeal (MP) or proximal interphalangeal (PIP) joint of affected finger;
2. Prescribed by or in consultation with a healthcare provider experienced in injection procedures of the hand and in the treatment of Dupuytren's contracture;
3. Age  $\geq 18$  years;
4. Member has not received surgical treatment (e.g., fasciectomy, fasciotomy) on the selected primary joint within the last 90 days;
5. If two injections (two vials) are requested, they are for one of the following (a or b):
  - a. One cord affecting two joints in the same finger;
  - b. Two cords affecting two joints in the same hand;
6. Dose does not exceed 0.58 mg per injection (one vial per injection).

**Approval duration: 3 months (up to 2 injections)**

##### **B. Peyronie's Disease** (must meet all):

1. Diagnosis of Peyronie's disease with both of the following (a and b):

**CLINICAL POLICY****Collagenase Clostridium Histolyticum**

- a. Palpable plaque;
  - b. Curvature deformity of  $\geq 30$  degrees at the start of therapy;
  2. Prescribed by or in consultation with a healthcare provider experienced in the treatment of male urological diseases;
  3. Age  $\geq 18$  years;
  4. Dose does not exceed 0.58 mg per injection (one vial per injection).
- Approval duration: 3 months (up to 2 injections)**

**C. Other diagnoses/indications**

1. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

**II. Continued Therapy****A. Dupuytren's Contracture (must meet all):**

1. Previously received medication via Centene benefit or member has previously met initial approval criteria;
2. Last treatment was  $\geq 4$  weeks ago;
3. Request is for one or both of the following:
  - a. MP or PIP contracture remains in affected cord since previous injection and the contracture is  $> 5$  degrees;
  - b. A different MP or PIP contracture will be injected that has a positive table top test and a flexion contracture of  $\geq 20$  degrees;
4. If two injections (two vials) are requested, the injections are for one cord affecting two joints in the same finger or two cords affecting two joints in the same hand;
5. Member has not received  $\geq 3$  total injections per affected cord;
6. Member has not received surgical treatment (e.g., fasciectomy, fasciotomy) on the selected primary joint within the last 90 days;
7. If request is for a dose increase, new dose does not exceed 0.58 mg per injection (one vial per injection).

**Approval duration: 3 months (up to 2 injections)**

**B. Peyronie's Disease (must meet all):**

1. Previously received medication via Centene benefit or member has previously met initial approval criteria;
2. Documented curvature deformity of  $\geq 15$  degrees remaining since last treatment cycle;
3. Last treatment cycle was  $\geq 6$  weeks ago;
4. Member has received  $< 4$  treatment cycles (i.e.  $< 8$  injections [2 injections per cycle]);
5. If request is for a dose increase, new dose does not exceed 0.58 mg per injection (one vial per injection).

**Approval duration: 3 months (up to 2 injections)**

**C. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

**CLINICAL POLICY**

**Collagenase Clostridium Histolyticum**

- Approval duration: Duration of request or 6 months (whichever is less);** or
2. Refer to CP.PHAR.57 if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized).

**III. Diagnoses/Indications for which coverage is NOT authorized:**

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PHAR.57 or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

MP: metacarpophalangeal joint

PIP: proximal interphalangeal joint

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Dupuytren's contracture	<p>0.58 mg per injection into a palpable cord with a contracture of a MP joint or a PIP joint</p> <p>Injections (0.58 mg) and finger extension procedures (24 hours later) may be administered up to 3 times per cord at approximately 4-week intervals. Up to 2 injections in the same hand may be performed during a treatment visit. Two palpable cords affecting 2 joints may be injected or 1 palpable cord affecting 2 joints in the same finger may be injected at 2 locations during a treatment visit. If a patient has other palpable cords with contractures of the MP or PIP joints, these cords may be injected at other treatment visits approximately 4 weeks apart.</p>	0.58 mg/dose
Peyronie's disease	<p>0.58 mg per injection administered into a Peyronie's plaque; if more than one plaque is present, inject into the plaque causing the curvature deformity.</p> <p>A treatment course consists of a maximum of 4 treatment cycles. Each treatment cycle consists of two Xiaflex injection procedures and one penile modeling procedure. The second Xiaflex injection procedure is performed 1 to 3 days after the first. The penile modeling procedure is performed 1 to 3 days after the second injection of the treatment cycle. The interval between treatment cycles is approximately six weeks. The treatment course therefore, consists of a maximum of 8 injection procedures and 4 modeling procedures.</p>	0.58 mg/dose

**CLINICAL POLICY**  
**Collagenase Clostridium Histolyticum**

	<p>If the curvature deformity is less than 15 degrees after the first, second or third treatment cycle, or if the healthcare provider determines that further treatment is not clinically indicated, then the subsequent treatment cycles should not be administered.</p> <p>The safety of more than one treatment course of Xiaflex is not known.</p>	
--	--	--

**VI. Product Availability**

Lyophilized powder for reconstitution (single-use glass vials): 0.9 mg of collagenase clostridium histolyticum

**VII. References**

1. Xiaflex Prescribing Information. Malvern, PA: Endo Pharmaceuticals, Inc.; July 2017. Available at <https://www.xiaflex.com/>. Accessed August 22, 2017.
2. Schulze SM and Tursi JP. Postapproval clinical experience in the treatment of Dupuytren’s contracture with collagenase clostridium histolyticum (CCH): the first 1,000 days. *Hand*. 2014; 9: 447-458.
3. Collagenase Drug Monograph. Clinical Pharmacology. Accessed August 2017. <http://www.clinicalpharmacology-ip.com>.

**Coding Implications**

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J0775	Injection, collagenase, clostridium histolyticum, 0.01 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Converted embedded SGM document into Centene clinical policy template	08.13	
Added treatment of Peyronie’s disease	12.13	01.14
Figure 1 :Removed the requirement that Xiaflex must be ordered through the REMS program Added the requirement for positive table top test Removed requirement that patient must not be a surgical candidate Figure 3: Removed the requirement that Xiaflex must be ordered through the REMS program; Added Appendix with definition of table top test	12.14	01.15

**CLINICAL POLICY**  
Collagenase Clostridium Histolyticum

Reviews, Revisions, and Approvals	Date	P&T Approval Date
<p>Clinical Information: Added information about Xiaflex managed distribution program and REMS program for both FDA approved indication</p> <p>Figure 2: rephrased question on how many injections received and clarified if contracture still remains rather than a reduction in contracture</p>		
<p>Converted policy to new template.</p> <p>DC initial auth criteria: added question about provider specialty per PI; removed question about underlying neuromuscular disorder affecting the hands - not in PI; removed thumb exclusion; added contraindications; changed approval from one injection to up to two total injections (same hand) visit.</p> <p>DC re-auth criteria: changed re-authorization approval of one injection to up to two injections but limited to not more than 3 injections in any one cord per PI; added criteria for a different contracture to be treated.</p> <p>PD criteria: added question about provider specialty and training per PI; added contraindications.</p>	12.15	01.16
<p>Converted policy to new template.</p> <p>Removed age limitation and added max dose for both indications.</p>	12.16	01.17
<p>Converted to new template.</p> <p>Dupuytren's and Peyronie's: Added age restriction as safety and effectiveness of Xiaflex in pediatric patients &lt; 18 years old have not been established and removed requirement related to history of hypersensitivity to Xiaflex per safety approach. Modified approval duration to allow up to 2 injections within a 3 month period.</p> <p>Peyronie's disease: Removed requirements related to completion of training for use of Xiaflex and contraindication related to plaques that involve the penile urethra since Xiaflex is available for the treatment of Peyronie's disease only through Xiaflex REMS program.</p>	08.22.17	11.17

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in

## CLINICAL POLICY

### Collagenase Clostridium Histolyticum

developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**CLINICAL POLICY****Collagenase Clostridium Histolyticum**

©2011 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.